



New entropy estimates for the Oldroyd-B model, and related models

Dan Hu, Tony Lelièvre

► To cite this version:

Dan Hu, Tony Lelièvre. New entropy estimates for the Oldroyd-B model, and related models. 2007.
hal-00135377

HAL Id: hal-00135377

<https://hal.science/hal-00135377>

Preprint submitted on 7 Mar 2007

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New entropy estimates for the Oldroyd-B model, and related models

D. Hu⁽¹⁾, T. Lelièvre^(2,3)

(1) School of Mathematical Sciences, Peking University, Beijing, 100871, P.R. China.

(2) CERMICS, Ecole Nationale des Ponts (ParisTech), 6 & 8 Av. B. Pascal, 77455 Marne-la-Vallée, France.

(3) INRIA Rocquencourt, MICMAC team, B.P. 105, 78153 Le Chesnay Cedex, France.
hdmxy@hotmail.com, lelievre@cermics.enpc.fr

7th March 2007

Abstract

This short note presents the derivation of a new *a priori* estimate for the Oldroyd-B model. Such an estimate may provide useful information when investigating the long-time behaviour of macro-macro models, and the stability of numerical schemes. We show how this estimate can be used as a guideline to derive new estimates for other macroscopic models, like the FENE-P model.

1 Introduction

We consider the Oldroyd-B model:

$$\text{Re} \left(\frac{\partial \mathbf{u}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{u} \right) = (1 - \varepsilon) \Delta \mathbf{u} - \nabla p + \text{div } \boldsymbol{\tau}, \quad (1)$$

$$\text{div } (\mathbf{u}) = 0, \quad (2)$$

$$\frac{\partial \boldsymbol{\tau}}{\partial t} + \mathbf{u} \cdot \nabla \boldsymbol{\tau} = \nabla \mathbf{u} \boldsymbol{\tau} + \boldsymbol{\tau} (\nabla \mathbf{u})^T - \frac{1}{\text{We}} \boldsymbol{\tau} + \frac{\varepsilon}{\text{We}} (\nabla \mathbf{u} + (\nabla \mathbf{u})^T), \quad (3)$$

where the Reynolds number $\text{Re} > 0$, the Weissenberg number $\text{We} > 0$ and $\varepsilon \in (0, 1)$ are some non-dimensional numbers. We suppose that the space variable \mathbf{x} lives in a bounded domain \mathcal{D} of \mathbb{R}^d . This system is supplied with initial conditions on the velocity \mathbf{u} and on the stress tensor $\boldsymbol{\tau}$. For simplicity, we assume no-slip boundary conditions on the velocity \mathbf{u} :

$$\mathbf{u} = 0 \text{ on } \partial \mathcal{D}. \quad (4)$$

We *suppose* that the initial data and the geometry are such that there exists a unique regular solution to (1)–(3) and our aim is to derive some *a priori* estimates on this solution.

Let us introduce the so-called conformation tensor $\mathbf{A} = \frac{\text{We}}{\varepsilon} \boldsymbol{\tau} + \text{Id}$. The partial differential equation (PDE) on $\boldsymbol{\tau}$ translates into the following PDE on \mathbf{A} :

$$\frac{\partial \mathbf{A}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{A} = \nabla \mathbf{u} \mathbf{A} + \mathbf{A} (\nabla \mathbf{u})^T - \frac{1}{\text{We}} \mathbf{A} + \frac{1}{\text{We}} \text{Id}. \quad (5)$$

One can check that if

$$\mathbf{A}(t=0) = \frac{\text{We}}{\varepsilon} \boldsymbol{\tau}(t=0) + \text{Id} \text{ is a positive definite symmetric matrix,} \quad (6)$$

then this property is propagated forward in time by (5) (and, in particular, $\boldsymbol{\tau}$ is symmetric). Assuming uniqueness of solution, this can be proven for example by

using the probabilistic interpretation of \mathbf{A} as a covariance matrix, as explained in Section 3. We will assume throughout this note that (6) is satisfied. Concerning the importance of positive-definiteness of \mathbf{A} , we refer for example to [7, Section 9.8.10] and also to the recent work [3, 4].

In Section 2, we recall how the classical *a priori* estimate for the Oldroyd-B model is derived. Next we show how it can be used to derive some bounds on the stress tensor, provided the initial condition satisfies $\det \mathbf{A}(t = 0) > 1$. In Section 3, we establish a new estimate, which comes from an entropy estimate on the micro-macro model associated with the Oldroyd-B model (see [5]). This estimate provides bounds on $(\mathbf{u}, \boldsymbol{\tau})$ without any assumption on $\boldsymbol{\tau}(t = 0)$ (apart from (6)). This new estimate could be useful to study the longtime behaviour of some macro-macro models, or to analyze the stability of some numerical schemes. Current research is directed towards clarifying this.

2 The classical estimate

Let us first introduce the kinetic energy:

$$E(t) = \frac{1}{2} \int_{\mathcal{D}} |\mathbf{u}|^2. \quad (7)$$

We easily obtain from (1)–(2):

$$\operatorname{Re} \frac{dE}{dt} = -(1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 - \int_{\mathcal{D}} \boldsymbol{\tau} : \nabla \mathbf{u}, \quad (8)$$

where for two matrices A and B , we denote $A : B = A_{i,j} B_{i,j} = \operatorname{tr}(A^T B)$. On the other hand, taking the trace of the PDE (3) on $\boldsymbol{\tau}$ and integrating over \mathcal{D} , we get:

$$\frac{d}{dt} \int_{\mathcal{D}} \operatorname{tr} \boldsymbol{\tau} = 2 \int_{\mathcal{D}} \nabla \mathbf{u} : \boldsymbol{\tau} - \frac{1}{\operatorname{We}} \int_{\mathcal{D}} \operatorname{tr} \boldsymbol{\tau}.$$

We thus obtain the following estimate:

$$\begin{aligned} & \frac{d}{dt} \left(\frac{\operatorname{Re}}{2} \int_{\mathcal{D}} |\mathbf{u}|^2 + \frac{1}{2} \int_{\mathcal{D}} \operatorname{tr} \boldsymbol{\tau} \right) \\ & + (1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 + \frac{1}{2\operatorname{We}} \int_{\mathcal{D}} \operatorname{tr} \boldsymbol{\tau} = 0. \end{aligned} \quad (9)$$

Remark 1 *In terms of \mathbf{A} , the energy estimate (9) writes:*

$$\begin{aligned} & \frac{d}{dt} \left(\frac{\operatorname{Re}}{2} \int_{\mathcal{D}} |\mathbf{u}|^2 + \frac{\varepsilon}{2\operatorname{We}} \int_{\mathcal{D}} \operatorname{tr} \mathbf{A} \right) \\ & + (1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 + \frac{\varepsilon}{2\operatorname{We}^2} \int_{\mathcal{D}} \operatorname{tr} (\mathbf{A} - \operatorname{Id}) = 0. \end{aligned} \quad (10)$$

In Lemma 1 below, we prove that $\operatorname{tr} \boldsymbol{\tau}$ is positive if $\det \mathbf{A}(t = 0) > 1$. This result combined with the estimate (9) thus yields some *a priori* bounds on $(\mathbf{u}, \boldsymbol{\tau})$ provided $\det(\mathbf{A})(t = 0) > 1$. In particular, it shows that \mathbf{u} and $\boldsymbol{\tau}$ go exponentially fast to 0 in the long time limit, using (9) and the Poincaré inequality: $\int_{\mathcal{D}} |\mathbf{u}|^2 \leq C \int_{\mathcal{D}} |\nabla \mathbf{u}|^2$.

Lemma 1 *Let us assume that $\det \mathbf{A}(t = 0) > 1$. Then, we have $\forall t \geq 0$, $\det \mathbf{A}(t) > 1$ and this implies that $\operatorname{tr} \boldsymbol{\tau}(t) > 0$.*

Proof: Using (5) and the Jacobi identity (which states that for any invertible matrix M depending smoothly on a parameter t , $\frac{d}{dt} \ln \det M = \operatorname{tr} (M^{-1} \frac{dM}{dt})$), we have:

$$\frac{\partial \ln(\det \mathbf{A})}{\partial t} + \mathbf{u} \cdot \nabla \ln(\det \mathbf{A}) = \frac{1}{\operatorname{We}} \operatorname{tr} (\mathbf{A}^{-1} - \operatorname{Id}). \quad (11)$$

Since for any symmetric positive matrix M of size $d \times d$,

$$(\det M)^{1/d} \leq (1/d) \operatorname{tr} M, \quad (12)$$

we obtain

$$\frac{\partial \ln(\det \mathbf{A})}{\partial t} + \mathbf{u} \cdot \nabla \ln(\det \mathbf{A}) \geq \frac{d}{\operatorname{We}} \left((\det \mathbf{A})^{-1/d} - 1 \right),$$

which we can rewrite in terms of $y = (\det \mathbf{A})^{1/d}$:

$$\operatorname{We} \left(\frac{\partial y}{\partial t} + \mathbf{u} \cdot \nabla y \right) \geq (1 - y). \quad (13)$$

This shows that $y > 1$ if $y(t=0) > 1$, and thus that $\det \mathbf{A} > 1$ if $\det \mathbf{A}(t=0) > 1$.

Indeed, using the characteristic method (by integrating the vector field $\mathbf{u}(t, \mathbf{x})$), one can rewrite (13) as

$$\operatorname{We} \frac{Dy}{Dt} \geq (1 - y).$$

Now, if y does not remain greater than 1, consider the first time t_0 such that $y(t_0) = 1$. We have on the one hand $\frac{Dy}{Dt}(t_0) < 0$ and, on the other hand $(1 - y(t_0)) = 0$. We reach a contradiction.

We thus have $\det \mathbf{A} > 1$ and therefore, using again (12), $\operatorname{tr} \boldsymbol{\tau} > d$. Since $\boldsymbol{\tau} = \frac{\varepsilon}{\operatorname{We}}(\mathbf{A} - \operatorname{Id})$, this is equivalent to $\operatorname{tr} \boldsymbol{\tau} > 0$. \diamond

Remark 2 If $\det \mathbf{A}(t=0) < 1$ (which is the case if $\operatorname{tr} \boldsymbol{\tau}(t=0) < 0$), Equation (13) shows that $\det \mathbf{A}$ grows along the characteristics as long as $\det \mathbf{A} < 1$.

3 Entropy estimate

We now consider a micro-macro (or multiscale) formulation of the Oldroyd-B model and some estimates based on entropy, inspired from [5].

3.1 General derivation of the entropy estimate for micro-macro models

We consider the following system:

$$\begin{cases} \operatorname{Re} \left(\frac{\partial \mathbf{u}}{\partial t}(t, \mathbf{x}) + \mathbf{u}(t, \mathbf{x}) \cdot \nabla \mathbf{u}(t, \mathbf{x}) \right) = (1 - \varepsilon) \Delta \mathbf{u}(t, \mathbf{x}) - \nabla p(t, \mathbf{x}) + \operatorname{div} \boldsymbol{\tau}(t, \mathbf{x}), \\ \operatorname{div} (\mathbf{u}(t, \mathbf{x})) = 0, \\ \boldsymbol{\tau}(t, \mathbf{x}) = \frac{\varepsilon}{\operatorname{We}} \left(\int_{\mathbb{R}^d} (\mathbf{X} \otimes \nabla \Pi(\mathbf{X})) \psi(t, \mathbf{x}, \mathbf{X}) d\mathbf{X} - \operatorname{Id} \right), \\ \frac{\partial \psi}{\partial t}(t, \mathbf{x}, \mathbf{X}) + \mathbf{u}(t, \mathbf{x}) \cdot \nabla_{\mathbf{x}} \psi(t, \mathbf{x}, \mathbf{X}) \\ = -\operatorname{div}_{\mathbf{X}} \left(\left(\nabla_{\mathbf{x}} \mathbf{u}(t, \mathbf{x}) \mathbf{X} - \frac{1}{2\operatorname{We}} \nabla \Pi(\mathbf{X}) \right) \psi(t, \mathbf{x}, \mathbf{X}) \right) + \frac{1}{2\operatorname{We}} \Delta_{\mathbf{X}} \psi(t, \mathbf{x}, \mathbf{X}). \end{cases} \quad (14)$$

This system is supplied with initial conditions on the velocity \mathbf{u} and on the distribution ψ . We recall that we suppose no-slip boundary conditions (4) on the velocity \mathbf{u} . This system corresponds to a micro-macro model of polymeric fluids, the polymer being modelled by two beads linked by a spring with potential energy Π . The configurational variable $\mathbf{X} \in \mathbb{R}^d$ models the end-to-end vector of the polymer. For more details on the modelling, we refer to [1, 8].

Notice that we could rewrite the former system as a system coupling a PDE and a stochastic differential equation (SDE), replacing the last two equations by:

$$\boldsymbol{\tau}(t, \mathbf{x}) = \frac{\varepsilon}{\operatorname{We}} \left(\mathbb{E} (\mathbf{X}_t(\mathbf{x}) \otimes \nabla \Pi(\mathbf{X}_t(\mathbf{x}))) - \operatorname{Id} \right), \quad (15)$$

$$\begin{aligned} d\mathbf{X}_t(\mathbf{x}) + \mathbf{u}(t, \mathbf{x}) \cdot \nabla_{\mathbf{x}} \mathbf{X}_t(\mathbf{x}) dt \\ = \left(\nabla_{\mathbf{x}} \mathbf{u}(t, \mathbf{x}) \mathbf{X}_t(\mathbf{x}) - \frac{1}{2\operatorname{We}} \nabla \Pi(\mathbf{X}_t(\mathbf{x})) \right) dt + \frac{1}{\sqrt{\operatorname{We}}} d\mathbf{W}_t. \end{aligned} \quad (16)$$

There, \mathbb{E} denotes the expectation, \mathbf{W}_t denotes a d -dimensional standard Brownian motion independent from the initial condition $(\mathbf{X}_0(\mathbf{x}))_{\mathbf{x} \in \mathcal{D}}$ which is such that, $\forall \mathbf{x} \in \mathcal{D}$, the law of $\mathbf{X}_0(\mathbf{x})$ is $\psi(0, \mathbf{x}, \mathbf{X}) d\mathbf{X}$.

Let us introduce the kinetic energy:

$$E(t) = \frac{1}{2} \int_{\mathcal{D}} |\mathbf{u}|^2. \quad (17)$$

We easily obtain:

$$\text{Re} \frac{dE}{dt} = -(1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 - \frac{\varepsilon}{\text{We}} \int_{\mathcal{D}} \int_{\mathbf{R}^d} (\mathbf{X} \otimes \nabla \Pi(\mathbf{X})) : \nabla \mathbf{u} \psi. \quad (18)$$

We now introduce the entropy of the system, namely:

$$\begin{aligned} H(t) &= \int_{\mathcal{D}} \int_{\mathbf{R}^d} \psi(t, \mathbf{x}, \mathbf{X}) \ln \left(\frac{\psi(t, \mathbf{x}, \mathbf{X})}{\psi_{\infty}(\mathbf{X})} \right), \\ &= \int_{\mathcal{D}} \int_{\mathbf{R}^d} \Pi(\mathbf{X}) \psi(t, \mathbf{x}, \mathbf{X}) + \int_{\mathcal{D}} \int_{\mathbf{R}^d} \psi(t, \mathbf{x}, \mathbf{X}) \ln(\psi(t, \mathbf{x}, \mathbf{X})) + C, \end{aligned} \quad (19)$$

with

$$\psi_{\infty}(\mathbf{X}) = \frac{\exp(-\Pi(\mathbf{X}))}{\int_{\mathbf{R}^d} \exp(-\Pi(\mathbf{X}))}, \quad (20)$$

and $C = \ln(\int_{\mathbf{R}^d} \exp(-\Pi(\mathbf{X})) d\mathbf{X})$. The function H is actually the relative entropy of ψ with respect to the equilibrium distribution ψ_{∞} .

After some computations (see [5]), we obtain:

$$\frac{dH}{dt} = -\frac{1}{2\text{We}} \int_{\mathcal{D}} \int_{\mathbf{R}^d} \psi \left| \nabla \ln \left(\frac{\psi}{\psi_{\infty}} \right) \right|^2 + \int_{\mathcal{D}} \int_{\mathbf{R}^d} (\mathbf{X} \otimes \nabla \Pi(\mathbf{X})) : \nabla \mathbf{u} \psi. \quad (21)$$

Therefore, introducing the free energy $F(t) = E(t) + \frac{\varepsilon}{\text{We}} H(t)$ of the system, we have:

$$\boxed{\begin{aligned} &\frac{d}{dt} \left(\frac{\text{Re}}{2} \int_{\mathcal{D}} |\mathbf{u}|^2 + \frac{\varepsilon}{\text{We}} \int_{\mathcal{D}} \int_{\mathbf{R}^d} \psi \ln \left(\frac{\psi}{\psi_{\infty}} \right) \right) \\ &+ (1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 + \frac{\varepsilon}{2\text{We}^2} \int_{\mathcal{D}} \int_{\mathbf{R}^d} \psi \left| \nabla \ln \left(\frac{\psi}{\psi_{\infty}} \right) \right|^2 = 0. \end{aligned}} \quad (22)$$

Using a logarithmic Sobolev inequality with respect to ψ_{∞} and a Poincaré inequality for $\mathbf{u} \in H_0^1(\mathcal{D})$, one can then obtain exponential convergence to equilibrium $\lim_{t \rightarrow \infty} (\mathbf{u}, \psi) = (0, \psi_{\infty})$ (see [5]). For some generalizations to the case $\mathbf{u} \neq 0$ on $\partial \mathcal{D}$, we refer to [5].

3.2 The Oldroyd-B case

Let us consider the Hookean dumbbell model, for which the potential Π of the entropic force is:

$$\Pi(\mathbf{X}) = \frac{\|\mathbf{X}\|^2}{2}. \quad (23)$$

By Itô's calculus, it is easy to derive from (16) that $\mathbf{A} = \mathbb{E}(\mathbf{X}_t \otimes \mathbf{X}_t)$ satisfies the following PDE:

$$\frac{\partial \mathbf{A}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{A} = \nabla \mathbf{u} \mathbf{A} + \mathbf{A} (\nabla \mathbf{u})^T - \frac{1}{\text{We}} \mathbf{A} + \frac{1}{\text{We}} \text{Id}. \quad (24)$$

This translates into the following PDE for $\boldsymbol{\tau} = \frac{\varepsilon}{\text{We}} (\mathbf{A} - \text{Id})$:

$$\frac{\partial \boldsymbol{\tau}}{\partial t} + \mathbf{u} \cdot \nabla \boldsymbol{\tau} = \nabla \mathbf{u} \boldsymbol{\tau} + \boldsymbol{\tau} (\nabla \mathbf{u})^T - \frac{1}{\text{We}} \boldsymbol{\tau} + \frac{\varepsilon}{\text{We}} (\nabla \mathbf{u} + (\nabla \mathbf{u})^T). \quad (25)$$

The Hookean dumbbell model is thus equivalent to the Oldroyd-B model (at least for regular enough solutions).

If $\psi(0, \mathbf{x}, \cdot)$ is Gaussian (with zero mean), so is $\psi(t, \mathbf{x}, \cdot)$:

$$\psi(t, \mathbf{x}, \mathbf{X}) = \frac{1}{(2\pi)^{d/2} \sqrt{\det(\mathbf{A})}} \exp\left(-\frac{\mathbf{X}^T \mathbf{A}^{-1} \mathbf{X}}{2}\right)$$

where $\mathbf{A} = \mathbb{E}(\mathbf{X}_t \otimes \mathbf{X}_t) = \int_{\mathbb{R}^d} \mathbf{X} \otimes \mathbf{X} \psi(t, \mathbf{x}, \mathbf{X}) d\mathbf{X}$ denotes as above the covariance matrix of \mathbf{X}_t , which depends on time and also on the space variable \mathbf{x} . The covariance matrix \mathbf{A} is symmetric and nonnegative. Moreover, since for almost all $t \geq 0$, $\int_{\mathcal{D}} \int_{\mathbb{R}^d} \psi(t, \mathbf{x}, \mathbf{X}) \ln\left(\frac{\psi(t, \mathbf{x}, \mathbf{X})}{\psi_\infty(\mathbf{X})}\right) < \infty$, then for almost all $t \geq 0$ and for almost all $\mathbf{x} \in \mathcal{D}$, \mathbf{A} is positive.

The following explicit expression of the relative entropy can then be derived:

$$\int_{\mathcal{D}} \int_{\mathbb{R}^d} \psi(t, \mathbf{x}, \mathbf{X}) \ln\left(\frac{\psi(t, \mathbf{x}, \mathbf{X})}{\psi_\infty(\mathbf{X})}\right) d\mathbf{X} = \int_{\mathcal{D}} \frac{1}{2} (-\ln(\det \mathbf{A}) - d + \text{tr} \mathbf{A}) .$$

On the other hand,

$$\int_{\mathcal{D}} \int_{\mathbb{R}^d} \psi(t, \mathbf{x}, \mathbf{X}) \left| \nabla_{\mathbf{X}} \ln\left(\frac{\psi(t, \mathbf{x}, \mathbf{X})}{\psi_\infty(\mathbf{X})}\right) \right|^2 d\mathbf{X} = \int_{\mathcal{D}} \text{tr}((\text{Id} - \mathbf{A}^{-1})^2 \mathbf{A}) .$$

Rewriting (22), we thus obtain the following estimate, in terms of \mathbf{A} :

$$\boxed{\begin{aligned} & \frac{d}{dt} \left(\frac{\text{Re}}{2} \int_{\mathcal{D}} |\mathbf{u}|^2 + \frac{\varepsilon}{2\text{We}} \int_{\mathcal{D}} (-\ln(\det \mathbf{A}) - d + \text{tr} \mathbf{A}) \right) \\ & + (1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 + \frac{\varepsilon}{2\text{We}^2} \int_{\mathcal{D}} \text{tr}((\text{Id} - \mathbf{A}^{-1})^2 \mathbf{A}) = 0. \end{aligned}} \quad (26)$$

This is, in the specific case of Hookean dumbbells (that is Oldroyd-B model) the macroscopic version of (22).

Since $-\ln(\det \mathbf{A}) - d + \text{tr}(\mathbf{A}) \geq 0$, this energy estimate yields some *a priori* bounds on (\mathbf{u}, \mathbf{A}) , and thus on $(\mathbf{u}, \boldsymbol{\tau})$. In sharp contrast to the classical estimate (9), it provides bounds on $(\mathbf{u}, \boldsymbol{\tau})$ without any assumption on $\boldsymbol{\tau}(t=0)$ (apart from (6)). Using a Poincaré inequality and the fact¹ that, for any symmetric positive matrix M of size $d \times d$,

$$-\ln(\det M) - d + \text{tr} M \leq \text{tr}((\text{Id} - M^{-1})^2 M)$$

exponential convergence to equilibrium ($\lim_{t \rightarrow \infty} (\mathbf{u}, \mathbf{A}) = (0, \text{Id})$) can be obtained from (26).

Remark 3 Notice that (26) can be schematically obtained as (10) $- \frac{\varepsilon}{2\text{We}} \int_{\mathcal{D}} (11)$.

Remark 4 If $\psi(0, \mathbf{x}, \cdot)$ is not Gaussian, it is always possible to replace it by a Gaussian initial condition with the same mean and variance, so that the macroscopic quantities $(\mathbf{u}, p, \mathbf{A})$ would be the same for the two initial conditions.

3.3 Application to related macroscopic models

The energy estimate (26) can be used as a guideline to derive energy estimates for other macroscopic models, even though they cannot be recast as a microscopic model of the form (14).

¹which can be seen as the logarithmic Sobolev inequality for Gaussian random variables translated on their covariance matrices

Let us consider the example of the FENE-P model [9, 2], for which

$$\boldsymbol{\tau} = \frac{\varepsilon}{\text{We}} \left(\frac{\mathbf{A}}{1 - \text{tr}(\mathbf{A})/b} - \text{Id} \right), \quad (27)$$

$$\frac{\partial \mathbf{A}}{\partial t} + \mathbf{u} \cdot \nabla \mathbf{A} = \nabla \mathbf{u} \mathbf{A} + \mathbf{A} (\nabla \mathbf{u})^T - \frac{1}{\text{We}} \frac{\mathbf{A}}{1 - \text{tr}(\mathbf{A})/b} + \frac{1}{\text{We}} \text{Id}. \quad (28)$$

For this model, we assume (6), and also that $\text{tr}(\mathbf{A})(t=0) < b$, and this property is propagated forward in time by (28) (see [6]).

Using the same ideas as for the Oldroyd-B model, we consider the “entropy” $H(t) = -\ln(\det \mathbf{A}) - b \ln(1 - \text{tr}(\mathbf{A})/b)$, and we compute its time-derivative:

$$\frac{d}{dt} \int_{\mathcal{D}} -b \ln(1 - \text{tr}(\mathbf{A})/b) = 2 \int_{\mathcal{D}} \frac{\nabla \mathbf{u} : \mathbf{A}}{1 - \text{tr}(\mathbf{A})/b} + \frac{1}{\text{We}} \int_{\mathcal{D}} \left(-\frac{\text{tr}(\mathbf{A})}{(1 - \text{tr}(\mathbf{A})/b)^2} + \frac{d}{1 - \text{tr}(\mathbf{A})/b} \right), \quad (29)$$

$$\frac{d}{dt} \int_{\mathcal{D}} \ln(\det(\mathbf{A})) = \frac{1}{\text{We}} \int_{\mathcal{D}} \left(-\frac{d}{1 - \text{tr}(\mathbf{A})/b} + \text{tr}(\mathbf{A}^{-1}) \right). \quad (30)$$

Combining these expressions with (8), we obtain

$$\boxed{\begin{aligned} & \frac{d}{dt} \left(\frac{\text{Re}}{2} \int_{\mathcal{D}} |\mathbf{u}|^2 + \frac{\varepsilon}{2\text{We}} \int_{\mathcal{D}} (-\ln(\det \mathbf{A}) - b \ln(1 - \text{tr}(\mathbf{A})/b)) \right) \\ & + (1 - \varepsilon) \int_{\mathcal{D}} |\nabla \mathbf{u}|^2 + \frac{\varepsilon}{2\text{We}^2} \int_{\mathcal{D}} \left(\frac{\text{tr}(\mathbf{A})}{(1 - \text{tr}(\mathbf{A})/b)^2} - \frac{2d}{1 - \text{tr}(\mathbf{A})/b} + \text{tr}(\mathbf{A}^{-1}) \right) = 0. \end{aligned}} \quad (31)$$

One can check that for any symmetric positive matrix M of size $d \times d$:

$$-\ln(\det(M)) - b \ln(1 - \text{tr}(M)/b) \geq -(b+d) \ln\left(\frac{b}{b+d}\right) \geq d \quad (32)$$

and that

$$-\ln(\det(M)) - b \ln(1 - \text{tr}(M)/b) + (b+d) \ln\left(\frac{b}{b+d}\right) \quad (33)$$

$$\leq \left(\frac{\text{tr}(M)}{(1 - \text{tr}(M)/b)^2} - \frac{2d}{1 - \text{tr}(M)/b} + \text{tr}(M^{-1}) \right). \quad (34)$$

The proof of these inequalities is tedious and can be done by diagonalizing the matrix M .

Equation (32) shows that

$$\frac{\text{Re}}{2} \int_{\mathcal{D}} |\mathbf{u}|^2 + \frac{\varepsilon}{2\text{We}} \int_{\mathcal{D}} \left(-\ln(\det \mathbf{A}) - b \ln(1 - \text{tr}(\mathbf{A})/b) + (b+d) \ln\left(\frac{b}{b+d}\right) \right)$$

is a non-negative quantity, and thus that (31) indeed yields some *a priori* bounds on (\mathbf{u}, \mathbf{A}) .

Equation (34) (which plays the role of the log-Sobolev inequality in the micro-macro models) shows that the estimate (31) can be used to prove exponential convergence to equilibrium.

References

- [1] R.B. Bird, C.F. Curtiss, R.C. Armstrong, and O. Hassager. *Dynamics of polymeric liquids*, volume 2. Wiley Interscience, 1987.
- [2] R.B. Bird, P.J. Dotson, and N.L. Johnson. Polymer solution rheology based on a finitely extensible bead-spring chain model. *J. Non-Newtonian Fluid Mech.*, 7:213–235, 1980. Errata: *J. Non-Newtonian Fluid Mech.*, 8 (1981) 193.

- [3] R. Fattal and R. Kupferman. Constitutive laws for the matrix-logarithm of the conformation tensor. *J. Non-Newtonian Fluid Mech.*, 123:281–285, 2004.
- [4] R. Fattal and R. Kupferman. Time-dependent simulation of viscoelastic flows at high Weissenberg number using the log-conformation representation. *J. Non-Newtonian Fluid Mech.*, 126:23–37, 2005.
- [5] B. Jourdain, C. Le Bris, T. Lelièvre, and F. Otto. Long-time asymptotics of a multiscale model for polymeric fluid flows. *Archive for Rational Mechanics and Analysis*, 181(1):97–148, 2006.
- [6] B. Jourdain and T. Lelièvre. Convergence of a stochastic particle approximation of the stress tensor for the FENE-P model, 2004. CERMICS 2004-263 report.
- [7] R. Keunings. *Fundamentals of Computer Modeling for Polymer Processing*, chapter Simulation of viscoelastic fluid flow, pages 402–470. Hanser, 1989.
- [8] H.C. Öttinger. *Stochastic Processes in Polymeric Fluids*. Springer, 1995.
- [9] A. Peterlin. Hydrodynamics of macromolecules in a velocity field with longitudinal gradient. *J. Polym. Sci. B*, 4:287–291, 1966.